

09/03/98

18:12:04

noy_mk4.asc

1

The budget and partitioning of reactive nitrogen species in the Arctic stratosphere

G.B. Osterman, B. Sen, G.C. Toon, R.J. Salawitch, J.J. Margitan, A.Y. Chang, and J.-F. Blavier

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

D.W. Fahey, R.S. Gao, L.A. Del Negro, S.G. Donnelly, E.R. Keim and J.A. Neuman

NOAA Aeronomy Laboratory, Boulder, CO

M. Loewenstein

NASA AMES Research Center, Moffett Field, CA

M.K.W. Ko and C. Scott

AER, Cambridge, MA

Volume mixing ratio profiles of numerous gases including NO, NO₂, HNO₃, HNO₄, ClNO₃, N₂O₅ and N₂O were measured remotely from 8 to 38 km by the JPL MkIV FTIR solar absorption spectrometer during balloon flights from Fairbanks, Alaska (64.8N, 147.6W) on May 8 and July 8, 1997. The observed ratio of NO_x (=NO+NO₂) to NO_y (total reactive nitrogen) is 10 to 30% greater than calculated by a steady state model using standard photochemistry and constrained by MkIV measurements of long lived precursors (e.g., H₂O, CH₄, CO and N₂O) and SAGE II aerosol surface area. The persistence of this discrepancy to 38 km altitude suggests that processes involving aerosols, such as the reduction of HNO₃ on the surface of soot particles, cannot be the sole explanation.

Calculations using a 35% decrease for the rate of NO₂+OH→HNO₃ (the dominant sink of NO_x in the Arctic summer stratosphere) compare well with MkIV observations of NO_x/NO_y and the individual NO_y species at all altitudes. The good agreement between theory and observation of N₂O₅ suggests that the heterogeneous hydrolysis of N₂O₅, a minor sink of NO_x during arctic summer, is handled correctly in the model. The MkIV NO_y vs N₂O relation agrees reasonably well with relations determined using the AER 2D model. In addition, comparison of the MkIV NO_y vs N₂O relation measured outside the vortex to ER-2 observations obtained inside the polar vortex on April 26, 1997 provides a context for inferring the degree of denitrification, which may have been as high as 5 ppbv during the 1996-97 Arctic winter.